

WHAT IS REDERMALISATION?

Succinic (amber) and hyaluronic acid are injected into the skin to stimulate skin regeneration and improve skin quality; signs of ageing are alleviated, the complexion is evened out and lymphatic drainage is accelerated.

Redermalisation works on three levels: it helps regulate biochemical processes in cells, maintains water balance and actively blocks free radicals.

Succinic acid normalizes microcirculation and accelerates lymphatic drainage, while also inhibiting the action of the tyrosinase enzyme and consequently maintaining an even skin tone. Hyaluronic acid activates anti-inflammatory mediators, restores water balance, contributes to the synthesis of basic structural proteins and provides mechanical support to the tissue.

The procedure is suitable for persons with facial swelling due to poor lymphatic drainage, under-eye bags, tired skin with fine lines and wrinkles, skin prone to hyperpigmentation and acne, where red and brown spots and scars remain.

It can be used on the skin of the face, neck and décolletage.

WHICH PRODUCTS DO WE USE?

Xela REDERM products are used.

HOW IS THE TREATMENT PERFORMED?

First, an anaesthetic cream is applied to make the procedure more comfortable. Then, a product of the appropriate composition is injected into the skin with a very thin needle, depending on the indication. After the procedure, a mask and a skincare product are applied.

HOW MANY REPETITIONS ARE NECESSARY?

The treatment is carried out every two weeks, three times in a row, and then repeated after six months.

THE SUCCESS OF THE PROCEDURE

After the first cycle of injections, the quality and tone of the skin improves. dark circles are less visible, the skin becomes lighter by one to two shades, and signs of fatigue disappear. Acne scars are less deep and the spots are lighter.

CONTRAINDICATIONS OR FOR WHOM IS THE PROCEDURE NOT SUITABLE?

The procedure should not be performed on pregnant women, breastfeeding mothers, children and persons with untreated epilepsy, chronic or acute inflammation of the skin at the injection site, known allergy to a component of the redermalisation product, serious organic and systemic diseases including autoimmune diseases, persons prone to hypertrophic scarring, and persons with a bleeding disorder or a tendency to haemorrhage. Special care is needed for people taking anti-clotting medication, as bleeding or bruising may occur at the site of the procedure. You should warn your doctor in case you are taking anti-clotting medication.

SIDE EFFECTS AND COMPLICATIONS

For a few days, you may see small bumps or lines at the injection site, which will disappear on their own. Light bleeding and transient bruising may be present.

POST-PROCEDURE INSTRUCTIONS

At home, a special care product is used three times a day, then the patient switches to own cosmetics. Any bruising can be covered with powder foundation the day after the procedure. Sauna is not recommended for 14 days.

You can return to work or your daily activities in one to three days.

Prior to the examination, you must also fill out the Informed Consent form together with the doctor as regulated by the Patient Rights Act and the Doctor's Duty to Obtain an Informed Consent. The rules for filling out the form and the statement are presented upon your arrival at our institution.

IF YOU HAVE ANY QUESTIONS REGARDING THIS BROCHURE, PLEASE CONTACT US FOR FURTHER INFORMATION.

We will take care of your health and well-being to the best of our ability. Thank you for your confidence in us.

Prepared by:

Janja Šmid Vauhnik, MD, Dermatovenerology Specialist